

Analyse of ionospheric and Geomagnetic Pre -earthquake Anomalies

Adibekyan M.V.

Ministry of Emergency Situations of the Republic of Armenia Territorial Survey for Seismic
Protection State Non-Commercial Organization
e-mail: adibekyan@yahoo.com

Abstract

Aiming at earthquake precursors apportionment the earthquake preparation displays of VARDENIS (Armenia, 29.04.2008, $M=3.7$) BORISAKHO (Georgia, 09.06.08, $M=4.1$), NAKHITCEVAN (Azerbaijan, 02.09.2008, $M=5.1$), earthquakes in time-series have been studied using the geomagnetic and ionosphere tools. Aiming at earthquake forecasting the anomaly in the ionosphere plasma is investigated by a radio-astronomical method. There were received some results, allowing to make out the difference of seismogenic anomalies of ionosphere between the longer anomalies connected to magnetic activity of ionosphere by the method of vertical reconnaissance of ionosphere.

Introduction

It is known, that the geophysical environment, including seismically active zones, is made up of solid, liquid and gaseous phases. It is known as well that in the zone of two phases separation a Zone of Separated Changes (ZSC) is forming, or as they are called in physical chemistry, double ionic (electric) layers. Depending on their structure, each of the ZSC of geophysical environment is characterized by capacity, inductivity and resistance (see [1]).

The results obtained earlier allow to make out the difference between activity of ionosphere, by the method of vertical reconnaissance of ionosphere.

Radio astronomical monitoring method makes possible, along with immediate detection of electromagnetic emission from the Earth's depth in the selected frequency band, to observe other types of lithospheric impact on the atmosphere, such as aerosol, electrostatic, acoustical/gravitational, etc. Those sources ultimately create anomalies at different altitudes of the atmosphere, high affect the propagating radio astronomical signal. Radio astronomical monitoring method has the following clear advantages over the active sensing methods: 1) Cosmic radio sources (Galactic background, discrete radio stars, etc.) are used instead of man-made radiation sources, often powerful, which are able to affect the observed layers of atmosphere; 2) Cosmic radio sources generate noise-like signals, which makes possible observation of the same source at several wavelengths (this is increasing the informative ness of monitoring); 3) Since the signals from the stellar radio sources pass through all layers of the atmosphere, their informative ness is high; 4) The radio astronomical instruments are highly sensitive and able to detect the smallest changes in the state of the atmosphere; 5) The method allows monitoring of solar activity by direct measurement of solar radiation density in the receiver waveband; 6) The cosmic source's power in the radio-frequencies range (over 10 MHz) is stable and known with high accuracy, while stability of the receiver system is provided by a controlled noise power source. This configuration allows reception of signals from point cosmic radio sources Swan and Cassiopeia-A, with nearly the same amplitudes of interference lobes. Time interval between these sources by the local meridian is 3 hours and 30 min.

The method and technique of research

A new Methodology has been elaborated that provides possibility to estimate the current Seismic hazard (its intensity, location and time) with a sufficiently big probability. The elaborated methodology was used for analysis of data received in the process of perpendicular

ionosphere from “Swan- A” and “Cassiopeia - A” point radio sources by radio astronomy methods. The time – series of geomagnetic field tension of T full vector.

Aiming at the earthquake forecasting the anomaly formations in the ionospheric plasma are investigated by a radio-astronomical method. For considering the geomagnetic field, the high accuracy proton magnetometers, which are measuring the T inductivity of geomagnetic field each 5 minutes are used for measuring the geomagnetic field. There have been used the time – series for Shooshi and Saravand ionospheric stations and Aruch and Bavra geomagnetic stations [2].




Fig.1. The time – series of the ionosphere field (Saravand station) obtained by Radio astronomical method for the VARDENIS (Armenia, 29.04.2008, $M=3.7$) earthquake.




Fig.2. The time – series of the geomagnetic field (Aruch, Bavra stations) for the VARDENIS (Armenia, 29.04.2008, $M=3.7$) earthquake.




Fig.3. The time – series of the ionosphere field (Shooshi station) obtained by Radio astronomical method for BORISAKHO (Georgia, 09.06.08, $M=4.1$) the earthquake.




Fig.4. The time – series of the ionosphere field (Saravand station) obtained by Radio astronomic method for the NAKHITCEVAN (Azerbaijan, 02.09.2008, $M=5.1$) earthquake.

Results

The results of the retrospective analysis of ionosphere observation data before NAKHITCEVAN (Azerbaijan, 02.09.2008, $M=5.1$), VARDENIS (Armenia, 29.04.2008, $M=3.7$) BORISAKHO (Georgia, 09.06.08, $M=4.1$), revealed the following basic types of anomaly (Figure. 15):

1. Blinking of ionosphere active radio-source Swan – A on the frequency of 74 MHz.
2. Anomaly of above – mentioned precursors is coming out up to 40 days before earthquake.

Conclusion

The results of analysis by used methods show that the anomalies generally appear on 1- 40 days before the earthquake.

References

- Badalyan M., Petrosyan. H., Adibekyan M.V. (2001) Peculiarities of Magnetic Precursors of Earthquakes on the Territory of Armenia. Journal of Georgian Geophysical Society. Physics of Solid Earth, vol.5, Tbilisi, p.37-43, 2001.
- Balassanian S. (1990), Dynamic Geo-electricity. Novosibirsk, “NAUKA”, Siberian Department, pp.232.